arXiv:1211.0444v1 [nucl-th] 2 Nov 2012

ABC EFFECT AS A SIGNAL OF CHIRAL SYMMETRY RESTORATION IN HADRONIC COLLISIONS

M. N. Platonova* and V. I. Kukulin†
Skobeltsyn Institute of Nuclear Physics,
Lomonosov Moscow State University,
Moscow 119991, Russia

A principally new mechanism for the basic 2π-fusion reaction $p + n \to d + (\pi\pi)_0$ in the energy region $T_p = 1.0$-1.4 GeV is suggested. The mechanism is aimed to provide a non-contradictory explanation for the comprehensive experimental studies of this reaction in exclusive setting done recently at COSY-WASA. The basic assumption of the model proposed is the production of the $I(J^P) = 0(3^+)$ dibaryon resonance D_{03} in the pn collision. The interference of two decay channels of this resonance: $D_{03} \to d + \sigma \to d + (\pi\pi)_0$ and $D_{03} \to D_{12} + \pi \to d + (\pi\pi)_0$ is shown to give a strong near-threshold enhancement in the $\pi\pi$ invariant mass spectrum, which is well known as the ABC-effect. The σ-meson parameters found to reproduce the ABC-enhancement are in a general agreement with models which predict a chiral symmetry restoration at a high excitation energy and/or high density of matter, although they are essentially less than those accepted for the free σ-meson. So, this result might be considered as an indication of the (partial) chiral symmetry restoration in dense and excited quark matter.

The famous ABC-effect discovered more than 50 years ago [1] is observed in double-pionic fusion reactions [1–3] as a pronounced spectral enhancement of isoscalar nature just above the $\pi\pi$ production threshold. The effect was initially interpreted [1] as being due to strong $\pi\pi$ rescattering in the scalar-isoscalar channel, associated naturally with the σ-meson. However the interpretation has been left since no narrow resonance with appropriate mass ($m \simeq 300$ MeV) was found in $\pi\pi$ scattering at low energies. At the same time, an other interpretation for the ABC-effect, based on the generation of two Δ-isobars via the t-channel meson exchange [4], was commonly accepted. Although the "t-channel Δ-Δ" mechanism did not provide quantitative description of the data, it allowed to reproduce the shape of differential cross sections found in the numerous inclusive experiments on double-pionic fusion [4, 5].

The situation has changed dramatically quite recently, after publication of the results of the first *exclusive and kinematically complete experiments* for the basic 2π-fusion reaction $p + n \to d + \pi^0\pi^0$ done at COSY-WASA [6]. The comparison of the new experimental data with theoretical predictions has demonstrated clearly that the above t-channel Δ-Δ model cannot reproduce even the qualitative behaviour of the experimental energy and angular distributions, giving just a low background in the considered energy region ($T_p = 1.0$-1.4 GeV). At the same time, the most intriguing discovery of these exclusive experiments was an observation of the well-pronounced resonance in the total 2π-production cross section. This fact has been interpreted by the authors [6] as a generation of the dibaryon resonance D_{03} in the pn collision, with quantum numbers $I(J^P) = 0(3^+)$, the mass $m_{D_{03}} \simeq 2.37$ GeV and the total width $\Gamma_{D_{03}} \simeq 70$ MeV. Such a resonance state has been predicted already in 1964 by Dyson and Xuong [7] and since then studied in numerous works, both theoretical [8–10] and experimental [11]. From the new exclusive experiments [6], the direct interrelation between the production and decay of the D_{03}-resonance and the ABC-effect has been clearly established. Having considered the D_{03} as the Δ-Δ bound state, the authors [6] suggested the virtual decay of the D_{03} into two Δ-isobars to be the basic mechanism for the reaction $p + n \to d + \pi^0\pi^0$. With such a "s-channel Δ-Δ" model they succeeded in a very good description of the numerous energy and angular distributions observed in this reaction. However, a reasonable agreement with the experimental data at low $\pi\pi$ invariant masses (in the region of the ABC-peak) could be reached in [6] only when using a very soft form factor $f_{\Delta\Delta}$ for the $D_{03} \to \Delta\Delta$ vertex with the cut-off parameter $\Lambda_{\Delta\Delta} = 0.15$ GeV/c. Such a low value of $\Lambda_{\Delta\Delta}$ means the characteristic radius of the D_{03}-state to be even larger than that of the deuteron. This is incompatible with the observed strong Δ-Δ binding in the D_{03}-state, $\epsilon_B(D_{03}) \simeq 90$ MeV, and also with the results of the various microscopic quark model calculations (see, e.g., [9]), which all predict the radius for $0(3^+)$ Δ-Δ bound state $r(D_{03}) \simeq 0.7$-0.9 fm, i.e. of the order of the nucleon one. Hence, the D_{03}-resonance appears to be the truly dibaryon state which arises in a situation when the quark cores of two Δ's are almost fully overlapped with each other. Moreover, the large width of the free Δ-isobar, $\Gamma_\Delta \simeq 120$ MeV, would not allow for two Δ's to go away to far distance, so the D_{03} system, even after the pion emission, is likely to stay in a dibaryon state with a small radius. So, this picture contradicts essentially to the concept of the bound state of two isolated quasi-free Δ-isobars, which therefore looks to be rather inconsistent. Thus, one is forced to consider some alternative models for the D_{03}-resonance.

---

*Electronic address: platonova@nucl-th.sinp.msu.ru
†Electronic address: kukulin@nucl-th.sinp.msu.ru

On the other hand, the isoscalar resonance D_{03} and its decay modes with the meson emission can be understood quite well within the framework of the dibaryon concept for short-range nuclear force [12]. In this concept, the conventional t-channel σ-meson exchange between two isolated nucleons is replaced by the s-channel σ exchange with the σ field surrounding the whole $6q$ bag, which appears in the overlap region of two nucleons. An emission of a light scalar meson occurs within a virtual transition of the $6q$ bag from the initial $2\hbar\omega$-excited quark configuration $|s^4p^2[42]\rangle$ to its ground state $|s^6[6]\rangle$. Then, a strong attraction of the σ field to the multi-quark core effectively induces a strong NN attraction at intermediate distances $r_{NN} \simeq 0.7$-0.8 fm (for the details, see [12] and references therein to the earlier works). The predictions of the model for the NN phase shifts as well as for the lightest nuclei properties [12, 13] are at the same level of accuracy as those of other modern NN-force models, like Bonn, Argonne, etc., still keeping quite moderate values for short-range cut-off parameters ($\Lambda_{\pi NN}$, etc.), which are compatible with the QCD and quark model estimations. However, within the dibaryon model, the best description of the experimental data has been achieved with the rather low mass of the σ-meson, $m_\sigma \simeq 350$ MeV, whereas in the conventional meson-exchange NN-force models the σ mass is taken to be ca. 600 MeV.

In fact, the mass and width of the σ-meson are still known with a large uncertainty: $m_\sigma = 400$-550 MeV, $\Gamma_\sigma = 400$-700 MeV [14]. This uncertainty is obviously related to the large width of the σ-meson, which makes it difficult to extract its parameters from experimental data. Another possible reason for the large spread of the σ-meson parameters may be the chiral symmetry restoration (CSR) effect which is known to occur in hadronic or nuclear media. The CSR in hot and/or dense nuclear matter has been studied in numerous works (see, e.g., [15, 16]). Besides that, it was shown [17] that the partial CSR takes place also in strongly excited states of *isolated hadrons* (baryons and mesons) at the excitation energies $E^* \gtrsim 500$ MeV. In fact, the rise of baryon density or nuclear matter temperature as well as a high hadron excitation energy lead to the increase of quark kinetic energy, which results in the suppression of the chiral condensate in QCD vacuum. This, in its turn, means the reduction of the σ-meson mass and the width for the $\sigma \to \pi\pi$ decay. As the degree of CSR depends on the conditions under which the σ-meson is produced, the σ masses and widths extracted from different experiments can also be different. Thus, since in the above dibaryon model the initial $6q$ bag is a dense object ($r_{6q} \simeq 0.5$-0.6 fm) and is also the $2\hbar\omega$-excited hadronic state, the renormalisation of the σ mass in the field of the $6q$ bag might be related to the partial CSR [12].

According to the above model [12], the deuteron wavefunction, besides the conventional NN component, has also the second, quark-meson component, which becomes dominating at short NN distances, i.e. when two nucleons are essentially overlapped with each other [18]. The second component of the deuteron has the structure $D_{01} \sim s^6 + \sigma \quad (l_\sigma = 0, 2)$ (a compact $6q$ bag dressed with a σ field), so it is similar in some sense to the picture of the physical nucleon in which the $3q$ core is dressed with the pionic cloud. Thus, analogously to the excited states of the nucleon, one can examine the excited states of the dibaryon D_{01} and classify them on their total angular momentum, isospin and parity. In this way, the experimentally observed D_{03} can be considered as a rotationally excited state of the D_{01}, with the quark-meson structure $s^6 + \sigma \quad (l_\sigma = 2, 4)$.

In fact, almost all dibaryon states lie in the vicinity of two-baryon thresholds, viz. NN, $N\Delta$, $\Delta\Delta$, etc., and are coupled strongly to the respective two-baryon channels. In our case, it is relevant to consider the following chain of dibaryon states with rising angular momenta: $D_{01} \sim NN$, $D_{12} \sim N\Delta$, $D_{03} \sim \Delta\Delta$, etc. Here the D_{12} is the well-known isovector ${}^1D_2$-dibaryon [19, 20] with quantum numbers $I(J^P) = 1(2^+)$ and the mass $m_{D_{12}} \simeq 2.15$ GeV. Although the D_{03} is a deeply bound state in the $\Delta\Delta$ channel, it is a resonance in the $p + n$ (as was observed in [6]) and $D_{12} + \pi$ systems. It becomes a resonance also in the $D_{01} + \sigma$ system, if the σ mass is less than 500 MeV. So, there are two basic possibilities for the decay of the D_{03}-resonance into the deuteron (i.e. into its quark-meson component D_{01}) and two pions:

(i) by an emission of the σ-meson (mainly in the d wave in respect to the $6q$ core due to the angular momentum conservation) which then decays into two pions;

(ii) by a sequential emission of two pions (each in p-wave) through an intermediate isovector dibaryon resonance D_{12}.

It is indicative that the above two interfering mechanisms for the excited dibaryon decay $D_{03} \to d + \pi\pi$ can be confronted with quite similar two mechanisms for the Roper resonance (excited nucleon) decay $N^*(1440) \to N + \pi\pi$ [14]: $N^*(1440) \to N + (\pi\pi)^{s\text{-}wave}_{I=0}$ (σ) and $N^*(1440) \to \Delta + \pi$. Note that the model [21] based on the excitation of the Roper resonance and its subsequent decay via the above two channels was quite successfully applied to the reactions $N + N \to d + \pi\pi$ and $N + N \to N + N + \pi\pi$ at the energies $T_N < 1$ GeV.

One can also treat other channels for the decay $D_{03} \to d + \pi\pi$, i.e. via a simultaneous emission of two uncorrelated pions without formation of the σ-meson or a sequential emission of two pions through other intermediate isovector dibaryons, such as the $1(1^-)$ and $1(3^-)$ (corresponding to the ${}^3P_1$ and ${}^3F_3$ NN partial waves). However the D_{12}-resonance should dominate, since it is well known [19] to give the largest contribution to the πd scattering cross section in the considered energy region. In a complete theoretical picture the conventional t-channel Δ-Δ mechanism should also be taken into account as the main background process to the D_{03} production.

In the present study, however, we consider just the dominating resonance contributions to the reaction $p +$

$n \to d + (\pi\pi)_0$, which are related to the above D_{03} decay channels (i) and (ii):

(a) $p + n \to D_{03} \to d + \sigma$, $\sigma \to (\pi\pi)_0$;

(b) $p + n \to D_{03} \to D_{12} + \pi$, $D_{12} \to d + \pi$.

The diagrams of the respective processes are shown in Fig. 1.



FIG. 1: The leading resonance mechanisms for the reaction $p + n \to d + (\pi\pi)_0$. The 3-momenta in the c.m.s. of two particles are indicated between the respective lines.

The amplitude for the emission of two neutral pions in the reaction $p + n \to d + \pi^0\pi^0$ at the c.m.s. energy $E = \sqrt{s}$ is then given by a sum of two terms:

$$\mathcal{M}_{\mu_i\mu_f} = \mathcal{M}^{(D_{03})}_{\mu_i}\left(\mathcal{M}^{(\sigma)}_{\mu_i\mu_f} + \mathcal{M}^{(D_{12})}_{\mu_i\mu_f}\right), \tag{1}$$

where

$$\mathcal{M}^{(D_{03})}_{\mu_i} = \frac{m^2_{D_{03}}\sqrt{\Gamma^{(2)}_{D_{03}np}/p}}{E^2 - m^2_{D_{03}} + im_{D_{03}}\Gamma_{D_{03}}}\mathcal{J}^{(D_{03})}_{\mu_i\mu_i}(\hat{p}), \tag{2}$$

$$\mathcal{M}^{(\sigma)}_{\mu_i\mu_f} = \frac{m_\sigma\sqrt{\Gamma^{(2)}_{D_{03}d\sigma}/q}\sqrt{\Gamma^{(0)}_{\sigma\pi^0\pi^0}/k}}{M^2_{\pi\pi} - m^2_\sigma + im_\sigma\Gamma_\sigma}\mathcal{J}^{(\sigma)}_{\mu_i\mu_f}(\hat{q}), \tag{3}$$

$$\mathcal{M}^{(D_{12})}_{\mu_i\mu_f} = \frac{1}{\sqrt{2}}\left(\frac{m_{D_{12}}\sqrt{\Gamma^{(1)}_{D_{03}D_{12}\pi^0}/k_1}\sqrt{\Gamma^{(1)}_{D_{12}d\pi^0}/\lambda_1}}{M^2_{d\pi} - m^2_{D_{12}} + im_{D_{12}}\Gamma_{D_{12}}}\right. \times \left.\mathcal{J}^{(D_{12})}_{\mu_i\mu_f}(\hat{k}_1, \hat{\lambda}_1) + \left[\vec{k}_1, \vec{\lambda}_1 \to \vec{k}_2, \vec{\lambda}_2\right]\right). \tag{4}$$

The amplitude $\mathcal{M}^{(D_{12})}_{\mu_i\mu_f}$ for the above process (b) is symmetrized over two identical pions.

When taking into account only the dominating, i.e. the lowest, partial waves in vertices (indicated in superscripts of Γ's in Eqs. (2)-(4)), the spin-angular terms $\mathcal{J}^{(\sigma)}_{\mu_i\mu_f}$ and $\mathcal{J}^{(D_{12})}_{\mu_i\mu_f}$ [22] can be calculated using the standard scheme for the angular momenta coupling, by decomposing the total angular momentum as $J = J1 + L$, i.e. $3 = 1 + 2$ for the process (a) and $\{3 = 2 + 1, 2 = 1 + 1\}$ for the process (b). The factor $\mathcal{J}^{(D_{03})}_{\mu_i\mu_i}(\hat{p})$ comes from the vertex $np \to D_{03}$ and, with the initial momentum $\vec{p}$ directed along z-axis, gives just a constant C_{μ_i}.

With the amplitudes defined in Eqs. (1)-(4), the differential cross sections as functions of the invariant masses squared $M^2_{\pi\pi}$ and $M^2_{d\pi}$ are given by

$$\frac{d\sigma}{d\,(M^2_{\pi\pi})} = \frac{\rho^{(\pi\pi)}}{(4\pi)^5 pE}\iint d\Omega_q d\Omega_k \frac{1}{3}\sum_{\mu_i,\mu_f}|\mathcal{M}_{\mu_i\mu_f}|^2, \tag{5}$$

$$\frac{d\sigma}{d\,(M^2_{d\pi})} = \frac{\rho^{(d\pi)}}{(4\pi)^5 pE}\iint d\Omega_{k_1} d\Omega_{\lambda_1} \frac{1}{3}\sum_{\mu_i,\mu_f}|\mathcal{M}_{\mu_i\mu_f}|^2, \tag{6}$$

where $\rho^{(\pi\pi)} = qk/2EM_{\pi\pi}$ and $\rho^{(d\pi)} = k_1\lambda_1/2EM_{d\pi}$ are the Lorentz-invariant phase-space factors.

The energy dependence for the partial width of the resonance R with the invariant mass M decaying into particles 1 and 2 with invariant masses M_1 and M_2 and the relative orbital angular momentum l has been parameterized as

$$\Gamma^{(l)}_{R12}(q) = \Gamma^{(l)*}_{R12}\left(\frac{q}{q^*}\right)^{2l+1}\left(\frac{(q^*)^2 + \varkappa^2}{q^2 + \varkappa^2}\right)^{l+1}, \tag{7}$$

where $q = \left[(M^2 - M_1^2 - M_2^2)^2 - 4M_1^2M_2^2\right]^{1/2}/2M$ is the modulus of the relative momentum between particles 1 and 2, and an asterisk denotes the values in the resonance point. Such a parametrization provides a correct near-threshold behaviour of the partial widths, however preventing an unphysical rise of the widths at high energies (see [23] for a similar parametrization in case $l = 1$). Thus, with an appropriate value of the parameter $\varkappa$, the centre of the Breit-Wigner distribution can be properly reproduced. For the partial widths introduced in Eqs. (2)-(4), this is achieved with $\varkappa = 0.1$-0.2 GeV/c.

The masses and total widths of the dibaryon resonances D_{03} and D_{12} have been fixed in our calculations as [24]

$$m_{D_{03}} = 2370 \text{ MeV}, \; \Gamma_{D_{03}} = 70 \text{ MeV},$$
$$m_{D_{12}} = 2150 \text{ MeV}, \; \Gamma_{D_{12}} = 110 \text{ MeV}.$$

The remaining model parameters, i.e. the mass and width of the σ-meson and the relative weight of the amplitudes corresponding to the processes (a) and (b), were derived from the fit to the experimental data [6] on the $M^2_{\pi\pi}$ spectrum, and then the $M^2_{d\pi}$ spectrum was calculated using the same parameter values.

The results for the $M^2_{\pi\pi}$ and $M^2_{d\pi}$ distributions at the peak energy $\sqrt{s} = 2.38$ GeV (where the total cross section has a maximum) are presented in Fig. 2. The results are normalised to the experimental value of the total cross section $(\sigma_T)_{\rm peak} \simeq 0.43$ mb. It is evident that our simple model reproduces the shapes of these two distributions almost perfectly. We observe that, although the cross section for the σ-generation process (a) alone is rather moderate, its contribution is crucial to reproduce the shape of the $M^2_{\pi\pi}$ distribution. Thus, its constructive interference with the D_{12}-production mechanism (b) at low $M^2_{\pi\pi}$ leads just to the observed height of the ABC-peak. On the other hand, when considering the $M^2_{d\pi}$ distribution (see Fig. 2b), the σ-production mechanism plays a role of a smooth background, while the process



FIG. 2: (Color online) Differential cross sections as functions of the invariant masses squared (a) $M^2_{\pi\pi}$ and (b) $M^2_{d\pi}$ in the reaction $pn \to d + \pi^0\pi^0$ at the energy $\sqrt{s} = 2.38$ GeV. The contribution of the σ-production mechanism (see Fig. 1a) is shown by dashed lines while the contribution of the mechanism going through the intermediate dibaryon D_{12} (see Fig. 1b) is shown by dash-dotted lines. The solid lines correspond to the summed cross sections. Shaded areas show the pure phase-space distributions. The experimental data (full circles) are taken from Ref. [6].

(b) alone almost gives the observed resonance enhancement.

The resonance peak in the $M^2_{d\pi}$ spectrum was associated previously with an excitation of the intermediate Δ-isobar [6]. However, our results show that this peak may reflect just a generation of the intermediate dibaryon resonance D_{12} which then decays into the final deuteron and pion. In fact, the suggested mechanism of two-pion emission through the D_{12} excitation is very similar to the s-channel Δ-Δ model proposed in [6], however without a soft form factor $f_{\Delta\Delta}$. The point is that the D_{12}-dibaryon, being located near to the $N\Delta$-threshold, has a large probability to be in the $N + \Delta$ $({}^5S_2)$ state. So, one needs additional tests to distinguish between these two mechanisms.

We also calculated the angular distributions for the final deuteron and pion emissions in the overall c.m. frame and then compared our model predictions with the experimental data. Our results for the angular distributions are shown in Fig. 3. The agreement with the data is not as good as for the invariant mass distributions, however it is still quite reasonable. Moreover, if we confront our model predictions with those found in [6] on the basis of the s-channel Δ-Δ model, the description of the above two angular distributions seems to be not worse than that reached in [6]. So, with only three basic parameters extracted from the $M^2_{\pi\pi}$ spectrum, our model is able to reproduce four differential distributions.



FIG. 3: (Color online) Angular distributions for the deuteron (a) and the pion (b) in the overall c.m.s. at the energy $\sqrt{s} = 2.38$ GeV. The meaning of curves is the same as in Fig. 2. The experimental data (full circles) are taken from Ref. [6].

The mass and width of the σ-meson extracted from the fit to the ABC-peak are

$$m_\sigma \simeq 300 \text{ MeV}, \; \Gamma_\sigma \simeq 90 \text{ MeV}.$$

These values are notably less than those quoted in PDG tables [14] for the σ-meson, and they are also lower than the values for the free σ mass and width, found by extrapolation from the dispersion relations for the $\pi\pi$ scattering amplitude to the σ complex pole [25],

$$m_\sigma^{(0)} = 441^{+16}_{-8} \text{ MeV}, \; \Gamma_\sigma^{(0)} = 544^{+18}_{-25} \text{ MeV}.$$

Bearing in mind the above arguments concerning the situations when the CSR occurs, one can suggest that the low values for the σ-meson parameters found here indicate to a partial CSR in the excited dibaryon state. Actually, the D_{03}-resonance represents dense quark matter (the density of a $6q$ system with a radius $r \simeq 0.8$ fm corresponds to about six-fold normal nuclear density) and has an excitation energy of 500 MeV above the deuteron pole. Thus, the σ-meson produced from the D_{03} decay should have the lower mass and width than those for the free σ-meson. As the σ width found here is still quite large, the σ-meson is likely to decay *before* it escapes the field of the multi-quark bag and esquires its free-space parameters. This implies that when measuring the $\pi\pi$ invariant mass distribution, one should observe just the renormalised σ-meson with the reduced mass and width. This conclusion is in an agreement with the results of numerous theoretical studies concerning the CSR in hadronic and nuclear media [15–17]. The further experimental and theoretical efforts are called for to check the fundamental CSR effects in hadronic systems.

To summarize, we have proposed a principally new model for the basic double-pionic fusion reaction $p+n \to d + (\pi\pi)_0$. The model takes into account the D_{03}-dibaryon production and its decay into the final deuteron and two pions by two alternative ways: (i) through an emission of the σ-meson and (ii) through a generation of the intermediate isovector dibaryon resonance D_{12}. So, the suggested mechanisms for the D_{03} decay have a remarkable reminiscence with two analogous modes of the Roper resonance $N^*(1440)$ decay. A reasonable agreement with the data of the recent exclusive experiments done at COSY-WASA [6], *without* an assumption of the unnaturally soft form factor in the vertex $D_{03} \to \Delta\Delta$, is obtained.

Within the model proposed, the ABC-peak is considered as a result of the σ-meson emission, whose mass and width, due to the partial restoration of chiral symmetry, are reduced in the field of the multi-quark bag as compared to their free-space values. In this way, the observed enhancement in the low $M_{\pi\pi}$ spectrum, similarly to the instant photograph, shows just the renormalised σ-meson in the field of the bag. Hence, by extracting the σ mass and width from the experimental data in the ABC region, one is able to judge about the degree of chiral symmetry restoration in the excited and/or dense hadronic systems. With this interpretation, it is easily understood why the ABC-effect is not seen in the reaction $pn \to pp\pi^-\pi^0$ [26]: although the D_{03}-resonance is produced there as well, but *the σ-meson is not.* So, we partially rehabilitate the initial interpretation of the ABC-effect suggested by its authors [1], even though the σ-meson generation in our model is not related to the $\pi\pi$ final state interaction. Thus, on the basis of the model proposed, one can treat the ABC-type experiments as a *first direct observation for the σ-meson production* in the NN, Nd, etc., collisions.

The authors are grateful to Prof. H. Clement, Drs. M. Bashkanov and T. Skorodko from Physical Institute of Tuebingen University for fruitful discussions on the COSY-WASA experimental results. One of the authors (V.I.K.) also thanks Prof. A. Faessler for the hospitality at Tuebingen University where this work was begun. The work was done under partial financial support from RFBR grants Nos. 10-02-00096 and 12-02-00908.

---

[1] A. Abashian, N.E. Booth, and K.M. Crowe, Phys. Rev. Lett. **5**, 258 (1960); **7**, 35 (1961).

[2] F. Plouin *et. al.*, Nucl. Phys. **A302**, 413 (1978).

[3] J. Banaigs *et. al.*, Nucl. Phys. **B105**, 52 (1976).

[4] T. Risser and M.D. Shuster, Phys. Lett. B **43**, 68 (1973); I. Bar-Nir, T. Risser, and M.D. Shuster, Nucl. Phys. **B87**, 109 (1975).

[5] G.W. Barry, Nucl. Phys. **B85**, 239 (1975).

[6] P. Adlarson *et. al.*, Phys. Rev. Lett. **106**, 242302 (2011); see also M. Bashkanov *et. al.*, Phys. Rev. Lett. **102**, 052301 (2009).

[7] F.J. Dyson and N.-H. Xuong, Phys. Rev. Lett. **13**, 815 (1964).

[8] J.L. Ping, H.X. Huang, H.R. Pang, F. Wang, and C.W. Wong, Phys. Rev. C **79**, 024001 (2009); T. Goldman, K. Maltman, G.J. Stephenson Jr., K.E. Schmidt, and F. Wang, Phys. Rev. C **39**, 1889 (1989).

[9] X.Q. Yuan, Z.Y. Zhang, Y.W. Yu, and P.N. Shen, Phys. Rev. C **60**, 045203 (1999); C.W. Wong, Phys. Rev. C **60**, 038201 (1999).

[10] H. Garcilazo, F. Fernandez, A. Valcarce, and R.D. Mota, Phys. Rev. C **56**, 84 (1997).

[11] H. Ikeda *et. al.*, Nucl. Phys. **B172**, 509 (1980) and refs. therein.

[12] V.I. Kukulin *et al.*, Ann. Phys. (N.Y.) **325**, 1173 (2010); V.I. Kukulin, I.T. Obukhovsky, V.N. Pomerantsev, and A. Faessler, Int. J. Mod. Phys. E **11**, 1 (2002); J. Phys. G **27**, 1851 (2001).

[13] V.I. Kukulin, I.T. Obukhovsky, P. Grabmayr, and A. Faessler, Phys. Rev. C **74**, 064005 (2006); V.I. Kukulin, V.N. Pomerantsev, M. Kaskulov, and A. Faessler, J. Phys. G **30**, 287 (2004).

[14] J. Beringer *et. al.* (Particle Data Group), Phys. Rev. D **86**, 010001 (2012).

[15] T. Hatsuda, T. Kunihiro, and H. Shimizu, Phys. Rev. Lett. **82**, 2840 (1999); T. Hatsuda and T. Kunihiro, Phys. Rev. Lett. **55**, 158 (1985).

[16] M.K. Volkov, A.E. Radzhabov, and N.L. Russakovich, Phys. Atom. Nucl. **66**, 997 (2003); M.K. Volkov *et. al.*, Phys. Lett. B **424**, 235 (1998).

[17] L. Ya. Glozman, Phys. Lett. B **475**, 329 (2000).

[18] The quark-meson component gives a small contribution (ca. 2-3%) to the total deuteron wavefunction normalisation, so it should be visible only when probing the deuteron structure with high-momentum probes [12].

[19] C.H. Oh, R.A. Arndt, I.I. Strakovsky, and R.L. Workman, Phys. Rev. C **56**, 635 (1997) and refs. therein; R. Bhandari, R.A. Arndt, L.D. Roper, and B.J. VerWest, Phys. Rev. Lett. **46**, 1111 (1981).

[20] N. Hoshizaki, Prog. Theor. Phys. **89**, 251 (1993).

[21] L. Alvarez-Ruso, Phys. Lett. B **452**, 207 (1999); L. Alvarez-Ruso, E. Oset, E. Hernandez, Nucl. Phys. **A633**, 519 (1998); see also T. Skorodko *et. al.*, Phys. Lett. B **695**, 115 (2011) for the updated version of the model.

[22] The production of the dibaryon resonance with quantum numbers $I(J^P) = 0(3^+)$ can occur in the np triplet state only, so the indices μ_i and μ_f run over all possible projections of the total spin $S = 1$ in initial and final states.

[23] S. Huber and J. Aichelin, Nucl. Phys. **A573**, 587 (1994).

[24] The D_{03} mass and width are taken from Ref. [6], and the parameters chosen for the D_{12} are close to those found in Ref. [20].

[25] I. Caprini, G. Colangelo, and H. Leutwyler, Phys. Rev. Lett. **96**, 132001 (2006).

[26] M. Bashkanov, private communication; see also A. Pricking, talk at the 12th International Workshop on Meson Production, Properties and Interaction “MESON 2012”, May 31 - June 5, 2012, Krakow, Poland.